

References

Pau D'Arco

- 18053
- Park BS, Lee KG, Shibamoto T, et al. Antioxidant activity and characterization of volatile constituents of Taheebo (*Tabebuia impetiginosa* Martius ex DC). *J Agric Food Chem* 2003;51:295-300. [View abstract](#).
- 18054
- Koyama J, Morita I, Tagahara K, Hirai K. Cyclopentene dialdehydes from *Tabebuia impetiginosa*. *Phytochemistry* 2000;53:869-72. [View abstract](#).
- 18055
- Park BS, Kim JR, Lee SE, et al. Selective growth-inhibiting effects of compounds identified in *Tabebuia impetiginosa* inner bark on human intestinal bacteria. *J Agric Food Chem* 2005;53:1152-7. [View abstract](#).
- 18056
- Park BS, Lee HK, Lee SE, et al. Antibacterial activity of *Tabebuia impetiginosa* Martius ex DC (Taheebo) against *Helicobacter pylori*. *J Ethnopharmacol* 2006;105:255-62. [View abstract](#).
- 18057
- Gómez Castellanos JR, Prieto JM, Heinrich M. Red Lapacho (*Tabebuia impetiginosa*)--a global ethnopharmacological commodity? *J Ethnopharmacol* 2009;121:1-13. [View abstract](#).
- 18058
- de Melo JG, Santos AG, de Amorim EL, et al. Medicinal plants used as antitumor agents in Brazil: an ethnobotanical approach. *Evid Based Complement Alternat Med* 2011;2011:365359. Epub 2011 Mar 8. [View abstract](#).
- 18059
- Kiage-Mokua BN, Roos N, Schrezenmeir J. Lapacho Tea (*Tabebuia impetiginosa*) Extract Inhibits Pancreatic Lipase and Delays Postprandial Triglyceride Increase in Rats. *Phytother Res* 2012 Mar 17. doi: 10.1002/ptr.4659. [View abstract](#).
- 18060
- Lemos OA, Sanches JC, Silva IE, et al. Genotoxic effects of *Tabebuia impetiginosa* (Mart. Ex DC.) Standl. (Lamiales, Bignoniaceae) extract in Wistar rats. *Genet Mol Biol* 2012;35:498-502. [View abstract](#).
- 68314
- Guerra, Mde O., Mazoni, A. S., Brandao, M. A., and Peters, V. M. Toxicology of Lapachol in rats: embryolethality. *Braz.J Biol.* 2001;61(1):171-174. [View abstract](#).
- 68315
- Felicio, A. C., Chang, C. V., Brandao, M. A., Peters, V. M., and Guerra, Mde O. Fetal growth in rats treated with lapachol. *Contraception* 2002;66(4):289-293. [View abstract](#).
- 68317
- Pereira, E. M., Machado, Tde B., Leal, I. C., Jesus, D. M., Damaso, C. R., Pinto, A. V., Giambiagi-deMarval, M., Kuster, R. M., and Santos, K. R. *Tabebuia avellanedae* naphthoquinones: activity against methicillin-resistant staphylococcal strains, cytotoxic activity and in vivo dermal irritability analysis. *Ann.Clin.Microbiol.Antimicrob.* 2006;5:5. [View abstract](#).
- 68318
- Lee, J. I., Choi, D. Y., Chung, H. S., Seo, H. G., Woo, H. J., Choi, B. T., and Choi, Y. H. beta-Lapachone induces growth inhibition and apoptosis in bladder cancer cells by modulation of Bcl-2 family and activation of caspases. *Exp.Oncol.* 2006;28(1):30-35. [View abstract](#).
- 68319
- Son, D. J., Lim, Y., Park, Y. H., Chang, S. K., Yun, Y. P., Hong, J. T., Takeoka, G. R., Lee, K. G., Lee, S. E., Kim, M. R., Kim, J. H., and Park, B. S. Inhibitory effects of *Tabebuia impetiginosa* inner bark extract on platelet aggregation and vascular smooth muscle cell proliferation through suppressions of arachidonic acid liberation and ERK1/2 MAPK activation. *J Ethnopharmacol.* 11-3-2006;108(1):148-151. [View abstract](#).
- 68320
- Woo, H. J., Park, K. Y., Rhu, C. H., Lee, W. H., Choi, B. T., Kim, G. Y., Park, Y. M., and Choi, Y. H. Beta-lapachone, a quinone isolated from *Tabebuia avellanedae*, induces apoptosis in HepG2

hepatoma cell line through induction of Bax and activation of caspase. J Med Food 2006;9(2):161-168. [View abstract](#)

68321

Kung, H. N., Chien, C. L., Chau, G. Y., Don, M. J., Lu, K. S., and Chau, Y. P. Involvement of NO/cGMP signaling in the apoptotic and anti-angiogenic effects of beta-lapachone on endothelial cells in vitro. J Cell Physiol 2007;211(2):522-532. [View abstract](#)

68322

de Cassia da Silveira E Sa and de Oliveira, Guerra M. Reproductive toxicity of lapachol in adult male Wistar rats submitted to short-term treatment. Phytother.Res. 2007;21(7):658-662. [View abstract](#)

68325

Kim, S. O., Kwon, J. I., Jeong, Y. K., Kim, G. Y., Kim, N. D., and Choi, Y. H. Induction of Egr-1 is associated with anti-metastatic and anti-invasive ability of beta-lapachone in human hepatocarcinoma cells. Biosci Biotechnol Biochem 2007;71(9):2169-2176. [View abstract](#)

68326

Yamashita, M., Kaneko, M., Iida, A., Tokuda, H., and Nishimura, K. Stereoselective synthesis and cytotoxicity of a cancer chemopreventive naphthoquinone from Tabebuia avellaneda. Bioorg.Med Chem.Lett. 12-1-2007;17(23):6417-6420. [View abstract](#)

68327

Bohler, T., Nolting, J., Gurragchaa, P., Lupescu, A., Neumayer, H. H., Budde, K., Kamar, N., and Klupp, J. Tabebuia avellaneda extracts inhibit IL-2-independent T-lymphocyte activation and proliferation. Transpl.Immunol. 2008;18(4):319-323. [View abstract](#)

68328

Savage, R. E., Tyler, A. N., Miao, X. S., and Chan, T. C. Identification of a novel glucosylsulfate conjugate as a metabolite of 3,4-dihydro-2,2-dimethyl-2H-naphtho[1,2-b]pyran-5,6-dione (ARQ 501, beta-lapachone) in mammals. Drug Metab Dispos. 2008;36(4):753-758. [View abstract](#)

68330

Queiroz, M. L., Valadares, M. C., Torello, C. O., Ramos, A. L., Oliveira, A. B., Rocha, F. D., Arruda, V. A., and Accorci, W. R. Comparative studies of the effects of Tabebuia avellaneda bark extract and beta-lapachone on the hematopoietic response of tumour-bearing mice. J Ethnopharmacol. 5-8-2008;117(2):228-235. [View abstract](#)

68331

Twardowschy, A., Freitas, C. S., Baggio, C. H., Mayer, B., dos Santos, A. C., Pizzolatti, M. G., Zacarias, A. A., dos Santos, E. P., Otuki, M. F., and Marques, M. C. Antiulcerogenic activity of bark extract of Tabebuia avellaneda, Lorentz ex Griseb. J Ethnopharmacol. 8-13-2008;118(3):455-459. [View abstract](#)

68332

Byeon, S. E., Chung, J. Y., Lee, Y. G., Kim, B. H., Kim, K. H., and Cho, J. Y. In vitro and in vivo anti-inflammatory effects of taheebo, a water extract from the inner bark of Tabebuia avellaneda. J Ethnopharmacol. 9-2-2008;119(1):145-152. [View abstract](#)

68333

Kung, H. N., Yang, M. J., Chang, C. F., Chau, Y. P., and Lu, K. S. In vitro and in vivo wound healing-promoting activities of beta-lapachone. Am.J Physiol Cell Physiol 2008;295(4):C931-C943. [View abstract](#)

91939

Block JB, Serpick AA, Miller W, Wiernik PH. Early clinical studies with lapachol (NSC-11905). Cancer Chemother Rep 2. 1974;4(4):27-8. [View abstract](#)

91940

Guiraud P, Steiman R, Campos-Takaki GM, Seigle-Murandi F, Simeon de Buochberg M. Comparison of antibacterial and antifungal activities of lapachol and beta-lapachone. Planta Med. 1994;60(4):373-4. [View abstract](#)

91941

de Almeida ER, da Silva Filho AA, dos Santos ER, Lopes CA. Antiinflammatory action of lapachol. J Ethnopharmacol. 1990 ;29(2):239-41. [View abstract](#)

91942

Kreher B, Lotter H, Cordell GA, Wagner H. New Furanonaphthoquinones and other Constituents of Tabebuia avellanedae and their Immunomodulating Activities in vitro. Planta Med. 1988 ;54(6):562-3. [View abstract](#)

91943

Paes JB, Morais VM, Lima CR. Resistência natural de nove madeiras do semi-árido brasileiro a fungos causadores da podridão-mole. R. Árvore, 2005;29(3):365-71.

91944

Nepomuceno JC. Lapachol and its derivatives as potential drugs for cancer treatment. In: Plants and Crop - The Biology and Biotechnology Research, 1st ed. iConcept Press Ltd.. Retrieved from:

https://www.researchgate.net/profile/Julio_Nepomuceno/publication/268378689_Lapachol_and_its_derivatives_as_potential_drugs_for_cancer_treatment/links/5469c8640cf20dedaf103e1.pdf.

91945

Awang DVC, Dawson BA, Ethier J-C, et al. Naphthoquinone Constituents of Commercial Lapacho/Pau d'arco/Taheebo Products. J Herbs Spic Med Plants. 1995;2(4):27-43.

91946

Awang DVC. Commercial taheebo lacks active ingredient. Information Letter 726 Can Pharm J. 1991;121:323-26.