

References

Gentian

374

Neubauer N, Marz RW. Placebo-controlled, randomized, double-blind, clinical trial with Sinupret sugar coated tablets on the basis of a therapy with antibiotics and decongestant nasal drops in acute sinusitis. *Phytomedicine* 1994;1:177-81.

379

Marz RW, Ismail C, Popp MA. Action profile and efficacy of a herbal combination preparation for the treatment of sinusitis. *Wien Med Wochenschr* 1999;149:202-8. [View abstract](#).

4912

Electronic Code of Federal Regulations. Title 21. Part 182 -- Substances Generally Recognized As Safe. Available at:

<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=182>

13435

Zagler B, Zelger A, Salvatore C, et al. Dietary poisoning with *Veratrum album*--a report of two cases. *Wien Klin Wochenschr* 2005;117:106-8. [View abstract](#).

13436

Toriumi Y, Kakuda R, Kikuchi M, et al. New triterpenoids from *Gentiana lutea*. *Chem Pharm Bull (Tokyo)* 2003;51:89-91. [View abstract](#).

13437

Tomic M, Tovilovic G, Butorovic B, et al. Neuropharmacological evaluation of diethylether extract and xanthones of *Gentiana kochiana*. *Pharmacol Biochem Behav* 2005;81:535-42. [View abstract](#).

13438

Haraguchi H, Tanaka Y, Kabbash A, et al. Monoamine oxidase inhibitors from *Gentiana lutea*. *Phytochemistry* 2004;65:2255-60. [View abstract](#).

13439

Uncini Manganelli RE, Chericoni S, Baragatti B. Ethnopharmacobotany in Tuscany: plants used as antihypertensives. *Fitoterapia* 2000;71:S95-100. [View abstract](#).

13440

Chericoni S, Testai L, Calderone V, et al. The xanthones gentiacaulein and gentiakochianin are responsible for the vasodilator action of the roots of *Gentiana kochiana*. *Planta Med* 2003;69:770-2. [View abstract](#).

13441

Baragatti B, Calderone V, Testai L, et al. Vasodilator activity of crude methanolic extract of *Gentiana kokiana* Perr. et Song. (Gentianaceae). *J Ethnopharmacol* 2002;79:369-72. [View abstract](#).

31135

Garnier, R., Carlier, P., Hoffelt, J., and Savidan, A. [Acute dietary poisoning by white hellebore (*Veratrum album* L.). Clinical and analytical data. A propos of 5 cases]. *Ann.Med.Interne (Paris)* 1985;136(2):125-128. [View abstract](#).

51701

Rojas, A., Bah, M., Rojas, J. I., and Gutierrez, D. M. Smooth muscle relaxing activity of gentiopicroside isolated from *Gentiana spathacea*. *Planta Med* 2000;66(8):765-767. [View abstract](#).

51702

Ozturk, N., Baser, K. H., Aydin, S., Ozturk, Y., and Calis, I. Effects of *Gentiana lutea* ssp. *sympyandra* on the central nervous system in mice. *Phytother Res* 2002;16(7):627-631. [View abstract](#).

51703

Kakuda, R., Machida, K., Yaoita, Y., Kikuchi, M., and Kikuchi, M. Studies on the constituents of *Gentiana* species. II. A new triterpenoid, and (S)-(+) and (R)-(-)-gentiolactones from *Gentiana lutea*. *Chem Pharm Bull.(Tokyo)* 2003;51(7):885-887. [View abstract](#).

51704

Wang, C. H., Wang, Z. T., Bligh, S. W., White, K. N., and White, C. J. Pharmacokinetics and tissue distribution of gentiopicroside following oral and intravenous administration in mice. *Eur J Drug Metab Pharmacokinet.* 2004;29(3):199-203. [View abstract](#)

51705

Schmieder, A., Schwaiger, S., Csordas, A., Backovic, A., Messner, B., Wick, G., Stuppner, H., and Bernhard, D. Isogentisin—a novel compound for the prevention of smoking-caused endothelial injury. *Atherosclerosis* 2007;194(2):317-325. [View abstract](#)

51706

Wang, C. H., Cheng, X. M., Bligh, S. W., White, K. N., Branford-White, C. J., and Wang, Z. T. Pharmacokinetics and bioavailability of gentiopicroside from decoctions of Gentianae and Longdan Xiegan Tang after oral administration in rats--comparison with gentiopicroside alone. *J Pharm Biomed.Anal* 9-3-2007;44(5):1113-1117. [View abstract](#)

51707

Aberham, A., Schwaiger, S., Stuppner, H., and Ganzen, M. Quantitative analysis of iridoids, secoiridoids, xanthones and xanthone glycosides in Gentiana lutea L. roots by RP-HPLC and LC-MS. *J Pharm Biomed.Anal* 11-5-2007;45(3):437-442. [View abstract](#)

51708

Citova, I., Ganzen, M., Stuppner, H., and Solich, P. Determination of gentisin, isogentisin, and amarogenitin in Gentiana lutea L. by capillary electrophoresis. *J Sep.Sci* 2008;31(1):195-200. [View abstract](#)

51709

Amin, A. Ketoconazole-induced testicular damage in rats reduced by Gentiana extract. *Exp Toxicol Pathol.* 2008;59(6):377-384. [View abstract](#)

51710

Chen, L., Liu, J. C., Zhang, X. N., Guo, Y. Y., Xu, Z. H., Cao, W., Sun, X. L., Sun, W. J., and Zhao, M. G. Down-regulation of NR2B receptors partially contributes to analgesic effects of Gentiopicroside in persistent inflammatory pain. *Neuropharmacology* 2008;54(8):1175-1181. [View abstract](#)

51711

Suzuki, O., Katsumata, Y., and Oya, M. Inhibition of monoamine oxidase by isogentisin and its 3-O-glucoside. *Biochem Pharmacol* 1978;27(16):2075-2078. [View abstract](#)

51712

Mathew, A., Taranalli, A. D., and Torgal, S. S. Evaluation of anti-inflammatory and wound healing activity of Gentiana lutea rhizome extract in animals. *Pharm.Biol.* 2004;42:8-12.

51713

Borgia, M., Sepe, N., Borgia, R., and Ori-Bellometti, M. Pharmacological activity of a herb extract: a controlled clinical study. *Current Therapeutic Research* 1981;29:525-536.

51714

Bricout, J. Identification and concentration of the bitter constituents of the roots of Gentiana lutea L. *Phytochemistry* 1974;13:2819-2823.

51715

Sticher, O. and Meier, B. Quantitative determination of the bitter principles in the roots of Gentiana lutea and Gentiana purpurea with HPLC [high-pressure liquid chromatography]. *Planta Medica* 1980;40:55-67.

51716

Schultze, J. and Franz, C. [Formation of bitter phenolic substances and sugar during growth phase of Genitana lutea roots [gentian, flavoring and perfume plant]]. *Planta Medica* 1978;33:275.

51717

Takino, K., Koshioka, M., and Kawaguchi, M. Quantitative determination of bitter components in gentianaceous plants. Studies on the evaluation of crude drugs. *Planta Medica* 1980;38:344-350.

51718

Szucs, Z., Danos, B., and Nyiredy, S. Comparative analysis of the underground parts of Gentiana species by HPLC with diode-array and mass spectrometric detection. *Chromatographia* 2002;56:S19.

51719

VanHaelen, M. and VanHaelen-Fastre, R. Quantitative determination of biologically active constituents in medicinal plant crude extracts by thin-layer chromatography-densitometry. I. *Aesculus hippocastaneum* L., *Arctostaphylos uva-ursi* Spreng., *Fraxinus excelsior* L., *Gentiana lutea* L., *Glycyrhiza glabra* L., *Hamamelis virginiana* L., *Hypericum perforatum* L., *Olea europaea* L., *Salix alba* L., and *Silybum marianum* Gaertn. *J Chromatogr* 1983;281:263-271.

51720

Hayashi, T. and Yamagishi, T. Two xanthone glycosides from *Gentiana lutea*. *Phytochemistry* 1988;27:3696-3699.